

- 2263, 6648, 3090, 6825, 4213, 4223.]
Infrasubspecific name: *rufino* (3251).
quercinum Thomson, 1877 (*Odontomerus quercinus*).
Palearctic. [Type: ♀, Sweden, SWD-LND-ZI. O.d.: 7493. Ref.: 4341, 1802, 2869, 2871, 6681, 6683, 7348, 4117, 5783, 4119, 7628, 1441, 5012, 5013, 3098, 1394, 192, 459, 363, 7706, 242, 5863, 257, 259, 1542, 5871, 3946, 2462, 2263, 6648, 3090, 6825, 4208, 4213, 4020, 4223.] Misidentified *quercinum* of authors = *Odontomerus thomsoni* Clément, 1938.
liogaster Szépligeti, 1914 (*Odontomerus*). [Type: Hungary, HGR-BDP-TMA. O.d.: 7348. Ref.: 4117, 4119.]
simile Habermehl, 1920 (*Odontomerus similis*). [Type: Germany. O.d.: 2895.]
brunneiventris Telenga, 1930 (*Odontomerus brunneiventris*). [Type: ♀, Russia, RSA-SPT-ZI. O.d.: 7396. Ref.: 5012.]
rufiventris Holmgren, 1860 (*Odontomerus*). Palearctic. [Type: ♀, Sweden. O.d.: 3412. Ref.: 7493, 906, 7334, 1802, 7282, 6681, 6683, 2048, 7348, 7127, 4117, 4119, 1439, 2652, 4121, 1629, 5012, 2656, 5791, 3098, 1394, 192, 6804, 363, 1536, 4062, 257, 1542, 3946, 6564, 4186, 3090, 6825, 4208, 4213, 4020, 4223.]
var. *rufipes* Ulbricht, 1911 (*Odontomerus rufiventris* var.). Palearctic. Considered as not available (3556). [Type: ♀, Germany, GMN-KFD-NM. O.d.: 7889. Ref.: 4116, 4117, 4119, 1394, 7158, 3556.]
rufum Uchida, 1928 (*Odontomerus rufus*). Palearctic. [Type: ♀, Japan, JPN-SPR-HU. O.d.: 7775. Ref.: 7713, 5155, 7706.]
sierrae Townes, 1960. Nearctic. [Type: ♀, U.S.A., USA-WST-USNM. O.d.: 7713. Ref.: 1165.]
spinipes Gravenhorst, 1829 (*Odontomerus*). Palearctic. [Type: ♀, Poland, PLD-WCW-IZU. O.d.: 2636. Ref.: 1696, 3999, 8386, 4593, 1802, 6681, 2476, 4127, 3098, 1394, 3262, 3263, 6804, 363, 3137, 7706, 4062, 5862, 257, 266, 1542, 5874, 3946, 3090, 6825, 4020, 4021.]
melanarium Holmgren, 1860 (*Odontomerus melanarium*). [Type: ♀, Sweden, SWD-SHM-NR. O.d.: 3412. Ref.: 7493, 1802, 7282, 6681, 6683, 7887, 5792, 7250, 7348, 7127, 4986, 4119, 2652, 5556, 5001, 5012, 5791, 5559, 3905, 31, 5570, 7706.]
stejnegeri Cushman, 1924 (*Odontomerus*). Palearctic. [Type: ♀, Russia, USA-WST-USNM. O.d.: 1724. Ref.: 7713, 7706.]
strangaliae Rohwer, 1917 (*Odontomerus*). Nearctic. [Type: ♀, U.S.A., USA-WST-USNM. O.d.: 6269. Ref.: 6271, 1735, 1739, 7648, 2189, 7710, 7713, 1165.]
thomsoni Clément, 1938 (*Odontomerus*). Palearctic. [Type: ♀, GMN-MCN-ZS. O.d.: 1394. Ref.: 3189, 3193, 3127, 363, 250, 252, 257, 258, 1542, 3946, 6564, 6648, 3090, 6825, 4213, 3544, 4223.] See also *Odontomerus quercinus* Thomson, 1877.
vicinum Cresson, 1870 (*Odontomerus vicinum*). Nearctic. [Type: ♂, U.S.A., USA-PLD-ANSP. O.d.: 1652. Ref.: 1664, 1802, 6681, 1666, 794, 6271, 1735, 1739, 5657, 7648, 7277, 7710, 7713, 1165.]
Xorides Latreille, 1809. [Type-species: *Ichneumon indicatorius* Latreille, 1806. O.d.: 4631. Ref.: 4632, 2630, 2636, 5356, 8328, 951, 6172, 3410, 3412, 7374, 7377, 2318, 7493, 1664, 6651, 149, 1802, 6681, 6682, 5179, 6683, 7990, 6271, 1208, 7775, 5012, 1394, 7648, 7710, 7655, 3441, 363, 7713, 382, 5488, 385, 7706, 7715, 242, 7682, 257, 7716, 2796, 5949, 1542, 1165, 2447, 2780, 2475.] Misidentified *Xorides* of authors = *Deuteroxorides* Viereck, 1914 or *Neoxorides* Clément, 1938.
Xorides (*Cyanoxorides*) Cameron, 1903. [Type-species: *Cyanoxorides brookei* Cameron, 1903. O.d.: 1040. Ref.: 6681, 7990, 7713, 382, 385, 7682, 7716, 2447, 2448, 2780, 2475.]
Spiloxorides Cameron, 1903. [Type-species: *Spiloxorides ruficeps* Cameron, 1903. O.d.: 1040. Ref.: 6681, 7990, 1744.]
abaddon Morley, 1913 (*Xylonomus*). Oriental. [Type: ♀, India, UKD-LDN-NHM. O.d.: 5199. Ref.: 5318, 7713, 7719, 2780, 8232.]
antennalis Szépligeti, 1914 (*Cyanoxorides*). Oriental. [Type: ♀, Malaysia, HGR-BDP-TMA. O.d.: 7348. Ref.: 7713, 7719, 2780.]
australensis Szépligeti, 1914 (*Cyanoxorides*). Australian. [Type: ♀, Australia, HGR-BDP-TMA. O.d.: 7348. Ref.: 7719, 2447, 2448, 2780.]
brookei Cameron, 1903 (*Cyanoxorides*). Oriental. [Type: ♀, Malaysia, UKD-LDN-NHM. O.d.: 1040. Ref.: 6681, 7713, 7719, 2780.]
caeruleus Cameron, 1905 (*Cyanoxorides*). Oriental. [Type: ♀, Malaysia, UKD-LDN-NHM. O.d.: 1066, 1059. Ref.: 1059, 6681, 7348, 7713, 7719, 2780.]
confusus Baltazar, 1961. Oriental. [Type: ♀, Philippines, PLP-MNL-BPI. O.d.: 382. Ref.: 388, 2780.]
crudelis Turner, 1919 (*Xylonomus*). Australian. [Type: ♀, Australia, UKD-LDN-NHM. O.d.: 7753. Ref.: 7713, 7719, 2447, 2448, 2780.]
gloriosus Szépligeti, 1914 (*Cyanoxorides*). Ethiopian. [Type: ♀, Ethiopia, HGR-BDP-TMA. O.d.: 7348. Ref.: 7716.]
iodes Baltazar, 1961. Oriental. [Type: ♀, Philippines, USA-WST-USNM. O.d.: 382. Ref.: 388, 5135, 2780.]
maculipennis Smith, 1859 (*Glypta*). Australian. [Type: ♀, Indonesia, UKD-OFD-HOPE. O.d.: 7074. Ref.: 7081, 1802, 6681, 7713, 7719, 2780.]
philippinensis Baltazar, 1961. Oriental. [Type: ♀, Philippines, USA-GVL-AEI. O.d.: 382. Ref.: 388, 2780.]
propodeus Cushman, 1933 (*Spiloxorides propodeus*). Oriental. [Type: ♀, China, GMN-EBW-DEI. O.d.: 1744. Ref.: 7713, 7719, 1279, 1337, 2780.]
ruficeps Cameron, 1903 (*Spiloxorides*). Oriental. [Type: ♀, Malaysia, UKD-LDN-NHM. O.d.: 1040. Ref.: 6681, 7713, 382, 7719, 388, 5135, 2780.]
erythrocephalus Cameron, 1905 (*Spiloxorides*). [Type: ♂, Malaysia, UKD-LDN-NHM. O.d.: 1066, 1059. Ref.: 1059, 6681, 7719, 2780.]
rufipleuralis Cushman, 1933 (*Spiloxorides*). Oriental. [Type: ♀, China, GMN-EBW-DEI. O.d.: 1744. Ref.: 7713, 7719, 5123, 1279, 1337, 2780.]
smithi Schmiedeknecht, 1907 (*Cyanoxorides*). Australian. [New name for *Glypta fracticornis* Smith, 1863. O.d.: 6681. Ref.: 7713, 7719, 2780.]
fracticornis Smith, 1863 (*Glypta*). Invalidated by pre-1961 replacement. Preoccupied by *Xorides*

- fracticornis* (Smith, 1860). [Type: ♀, Indonesia, UKD-OFD-HOPE. O.d.: 7077. Ref.: 7078, 7081, 1802, 6681, 2780.]
- spectabilis* Tosquinet, 1903 (*Moansa*). Australian. [Type: ♀, Papua New Guinea, IITL-GNA-MCSN. O.d.: 7639. Ref.: 6681, 7713, 7719, 2780.]
- splendens* Brues, 1918 (*Cyanoxorides*). Australian. [Type: ♀, Solomon Islands, USA-CBG-MCZ. O.d.: 942. Ref.: 7719, 2780.]
- Xorides* (*Epixorides*)** Smith, 1862. [Type-species: *Epixorides chalybeator* Smith, 1862. O.d.: 7076. Ref.: 149, 1802, 6681, 7990, 7713, 382, 385, 7682, 2780, 2475.]
- Ahyborhyssa* Enderlein, 1919. [Type-species: *Epirhyssa annulicornis* Cameron, 1899. O.d.: 2136.]
- caerulescens* Morley, 1913 (*Xylonomus*). Oriental. [Type: ♂, Sri Lanka, UKD-LDN-NHM. O.d.: 5199. Ref.: 2066, 487, 5318, 7713, 7719, 2780.]
- chalybeator* Smith, 1862 (*Epixorides*). Australian. [Type: ♂, Indonesia, UKD-OFD-HOPE. O.d.: 7076. Ref.: 7081, 1802, 6681, 7713, 7719, 2780.]
- citrimaculatus* Wang & Gupta, 1995. Oriental. [Type: ♀, China, CNA-BJG-ASIZ. O.d.: 8232.]
- elizabethae* Bingham, 1898 (*Xylonomus*). Oriental. [Type: ♀, India, GMN-BLN-ZMHU. O.d.: 646. Ref.: 1802, 6681, 5199, 4962, 5318, 7713, 7719, 5122, 2780, 8232.]
- annulicornis* Cameron, 1899 (*Epirhyssa*). [Type: ♀, India, UKD-LDN-NHM. O.d.: 1017. Ref.: 1802, 6681, 5199, 2136, 7751, 5318, 7719, 2780.]
- blanditor* Tosquinet, 1903 (*Xylonomus*). [Type: ♀, Indonesia, BGM-BXL-IRSN. O.d.: 7639. Ref.: 6681, 2780.]
- albolineatus* Cameron, 1912 (*Cyanoxorides*). [Type: ♀, Malaysia, UKD-LDN-NHM. O.d.: 1134. Ref.: 2780.]
- javanus* Szépligeti, 1914 (*Cyanoxorides*). [Type: ♀, Indonesia, HGR-BDP-TMA. O.d.: 7348. Ref.: 7719, 2780.]
- erythrothorax* Turner, 1919 (*Cyanoxorides*). Oriental. [Type: ♂, Laos, UKD-LDN-NHM. O.d.: 7751. Ref.: 7713, 7719, 2780.]
- flavipictus* Baltazar, 1961. Oriental. [Type: ♀, Philippines, USA-GVL-AEI. O.d.: 382. Ref.: 388, 2780.]
- formosanus* Sonan, 1936 (*Cyanoxorides*). Oriental. [Type: ♀, China, CNA-TPI-TARI. O.d.: 7149. Ref.: 7713, 7719, 1279, 1337, 7699, 2780.]
- fracticornis* Smith, 1860 (*Xylonomus*). Australian. [Type: ♀, Indonesia, UKD-OFD-HOPE. O.d.: 7075. Ref.: 7081, 1802, 6681, 7713, 7719, 2780.]
- fulgidipennis* Smith, 1858 (*Xylonomus*). Oriental. [Type: ♀, Malaysia, UKD-OFD-HOPE. O.d.: 7071. Ref.: 7081, 1802, 6681, 7713, 7719, 2780.]
- giganticus* Baltazar, 1961. Oriental. [Type: ♀, Philippines, USA-WST-USNM. O.d.: 382. Ref.: 388, 2780.]
- mindanensis* Baltazar, 1961. Oriental. [Type: ♀, Philippines, USA-WST-USNM. O.d.: 382. Ref.: 388, 5135, 2780, 8232.]
- nigriceruleus* Wang & Gupta, 1995. Oriental. [Type: ♀, China, CNA-BJG-ASIZ. O.d.: 8232.]
- rufomaculatus* Cameron, 1905 (*Cyanoxorides*). Oriental. [Type: ♀, Malaysia, UKD-LDN-NHM. O.d.: 1066, 1059. Ref.: 1059, 6681, 7348, 7713, 7719, 2780.]
- vitalisi* Turner, 1919 (*Cyanoxorides*). Oriental. [Type: ♀, Vietnam, UKD-LDN-NHM. O.d.: 7751. Ref.: 7713, 7719, 2780.]
- Xorides* (*Exomus*)** Townes, 1960. (*Xorides* subg.) [Type-species: *Anomalon humerale* Say, 1829. (*Anomalon humerale* sensu Townes, 1960.) O.d.: 7713. Ref.: 1330, 7682, 1165, 2475.]
- humeralis* Say, 1829 (*Anomalon humerale*). Nearctic, Neotropical. [Type: U.S.A. O.d.: 6584. Ref.: 6585, 1652, 6023, 6064, 6068, 1664, 2991, 1802, 6681, 7992, 6271, 1778, 1735, 7648, 2189, 7710, 7713, 5953, 1165, 6445.]
- australis* Cresson, 1870 (*Xylonomus*). [Type: ♀, U.S.A., USA-PLD-ANSP. O.d.: 1652. Ref.: 1655, 1664, 2991, 1802, 6681, 1666, 6271, 7648, 2189.]
- lavallensis* Provancher, 1874 (*Xylonomus*). [Type: ♀, Canada, CND-SFY-UL. O.d.: 6023. Ref.: 1802, 7648, 7713, 404.]
- piceatus* Rohwer, 1913 (*Xylonomus* (*Moerophora*)). [Type: ♀, U.S.A., USA-WST-USNM. O.d.: 6265. Ref.: 6271, 7710, 7713.]
- chiricensis* Townes, 1960 (*Xorides* (*Exomus*) *humerale* ssp.). [Type: ♀, U.S.A., USA-WST-USNM. O.d.: 7713.]
- excomptus* Townes, 1960 (*Xorides* (*Exomus*) *humerale* ssp.). [Type: ♀, U.S.A., USA-DVS-UC. O.d.: 7713.]
- mexicanus* Townes, 1960 (*Xorides* (*Exomus*) *humerale* ssp.). [Type: ♀, Mexico, USA-LRC-UK. O.d.: 7713.]
- sierrae* Townes, 1960 (*Xorides* (*Exomus*) *humerale* ssp.). [Type: ♀, U.S.A., USA-BKL-UC. O.d.: 7713.]
- Xorides* (*Gonioprymnus*)** Cameron, 1906. [Type-species: *Gonioprymnus maculiceps* Cameron, 1906. O.d.: 1081. Ref.: 6681, 2135, 7990, 510, 7713, 7682, 7716, 2475.]
- Lavaudenia* Seyrig, 1932. [Type-species: *Lavaudenia mirabilis* Seyrig, 1932. O.d.: 6885.]
- hulstaerti* Benoit, 1952 (*Gonioprymnus*). Ethiopian. [Type: ♂, Zaire, BGM-TVR-MRAC. O.d.: 510. Ref.: 7716.]
- maculatus* Benoit, 1952 (*Gonioprymnus*). Ethiopian. [Type: ♀, Zaire, BGM-TVR-MRAC. O.d.: 510. Ref.: 7716.]
- maculiceps* Cameron, 1906 (*Gonioprymnus*). Ethiopian. [Type: ♀, South Africa, SAF-CTN-SFM. O.d.: 1081. Ref.: 6681, 5212, 510, 7713, 7716.]
- unifasciatus* Morley, 1916 (*Xylonomus*). [Type: ♀, South Africa, SAF-CTN-SFM. O.d.: 5210. Ref.: 7716.]
- mirabilis* Seyrig, 1932 (*Lavaudenia*). Ethiopian. [Type: ♀, Madagascar, FCE-PRS-MNHN. O.d.: 6885. Ref.: 7713, 7716.]
- serratitibia* Enderlein, 1914 (*Gonioprymnus*). Ethiopian. [Type: ♀, Namibia, GMN-HBG-ZM. O.d.: 2135. Ref.: 7716.]
- tarsalis* Szépligeti, 1914 (*Gonioprymnus*). Ethiopian. [Type: ♀, Zaire, HGR-BDP-TMA. O.d.: 7348. Ref.: 7716.]
- Xorides* (*Gonophonus*)** Förster, 1869. [Type-species: *Gonophonus mokrzeckii* Kokujev, 1902.

- O.d.: 2318. Ref.: 6651, 149, 4147, 6682, 7990, 1394, 363, 7713, 382, 7719, 5701, 385, 7682, 1542, 2798, 2780, 2475.]
- Caenostoma* Cameron, 1905. [Type-species: *Caenostoma filicornis* Cameron, 1905. O.d.: 1066, 1061. Ref.: 6681, 7990, 382, 385.]
- Paraglypta* Kieffer, 1921. [Type-species: *Paraglypta tubigera* Kieffer, 1921. O.d.: 4085. Ref.: 7682.]
- annulator* Fabricius, 1804 (*Cryptus*). Palearctic. [Type: ♀, Austria, DMK-CPH-UZM. O.d.: 2174. Ref.: 2636, 7726, 1802, 6685, 8498, 7706, 242, 257, 1542, 3946, 6825, 4213, 4020, 4223, 4225.]
- circulator* Thunberg, 1822 (*Ichneumon*). Preoccupied by *Ichneumon circulator* Panzer, 1800. [Emendation for *Cryptus annulator* Fabricius, 1804. O.d.: 7537. Ref.: 7538.]
- fasciipennis* Kriechbaumer, 1894 (*Xylonomus*). [O.d.: 6637. Ref.: 1802, 6680, 6681, 7348, 1208, 4119, 5012, 1394, 1217, 1223, 192, 197, 363, 7713, 1239, 7706.]
- carinifrons* Baltazar, 1961. Oriental. [Type: ♀, Philippines, USA-GVL-AEI. O.d.: 382. Ref.: 388, 2798, 2780.]
- corcyrensis* Kriechbaumer, 1894 (*Xylonomus*). Palearctic. [O.d.: 6637. Ref.: 1802, 6681, 6683, 5012, 1394, 7713, 253, 257, 6899, 6898, 1542, 3946, 6825, 4213, 4223.]
- ssp. *persicator* Aubert, 1976 (*Xorides corcyrensis* ssp.). Palearctic. [Type: Iran. O.d.: 277.]
- exmacularis* Wang & Gupta, 1995. Palearctic. [Type: ♀, China, CNA-BJG-ASIZ. O.d.: 8232.]
- exquisitus* Tosquin, 1903 (*Moansa*). Oriental. [Type: ♀, Indonesia, BGM-BXL-IRSN. O.d.: 7639. Ref.: 6681, 7713, 382, 7719, 2798, 2780.]
- filicornis* Cameron, 1905 (*Caenostoma*). [Type: ♂, Malaysia, UKD-LDN-NHM. O.d.: 1066, 1061. Ref.: 1061, 6681, 2780.]
- tubiger* Kieffer, 1921 (*Paraglypta tubigera*). [Type: ♀, Vietnam. O.d.: 4085. Ref.: 2780.]
- ssp. *ceylonicus* Gupta & Chandra, 1977 (*Xorides* (*Gonophonus*) *exquisita* ssp.). Oriental. [Type: ♀, Sri Lanka, ATA-WEN-NM. O.d.: 2798. Ref.: 2780.]
- ssp. *xanthopleurus* Baltazar, 1961 (*Xorides* (*Caenostoma*) *exquisita* ssp.). Oriental. [Type: ♀, Philippines, USA-WST-USNM. O.d.: 382. Ref.: 388, 2798, 2780.]
- karnaticus* Gupta & Chandra, 1977. Oriental. [Type: ♀, India, USA-GVL-GUPTA. O.d.: 2798. Ref.: 2780.]
- propinquus* Tschek, 1869 (*Xylonomus*). Palearctic. [O.d.: 7737. Ref.: 1802, 6680, 6681, 4712, 1208, 5012, 1394, 2849, 5472, 1223, 192, 2864, 6804, 363, 1229, 7713, 1238, 242, 257, 1542, 3946, 6824, 6825, 4213, 4020, 2547, 4223.]
- mokrzechki* Kokujev, 1902 (*Gonophonus*). [Type: ♀, Ukraine. O.d.: 4147. Ref.: 5012.]
- scaber* Gravenhorst, 1829 (*Xylonomus*). Palearctic. [O.d.: 2636. Ref.: 1696, 1802, 6680, 6681, 6683, 6437, 5012, 1394, 363, 7713, 257, 1542, 3946, 6564, 4020.]
- Xorides* (*Gymnacanthus*) Benoit, 1952. [Type-species: *Gymnacanthus aciculatus* Benoit, 1952. O.d.: 510. Ref.: 7716, 2475.]
- aciculatus* Benoit, 1952 (*Gymnacanthus*). Ethiopian. [Type: ♀, Zaire, BGM-TVR-MRAC. O.d.: 510. Ref.: 7713, 7716.]
- mayumbensis* Benoit, 1952 (*Gymnacanthus*). Ethiopian. [Type: ♀, Zaire, BGM-TVR-MRAC. O.d.: 510. Ref.: 7716.]
- similis* Benoit, 1952 (*Gymnacanthus*). Ethiopian. [Type: ♀, Zaire, BGM-TVR-MRAC. O.d.: 510. Ref.: 7716.]
- Xorides* (*Moansa*) Tosquin, 1896. [Type-species: *Moansa praestans* Tosquin, 1896. O.d.: 7636. Ref.: 149, 1802, 6681, 7990, 5212, 7713, 7682, 7716, 2475.]
- praestans* Tosquin, 1896 (*Moansa*). Ethiopian. [Type: ♀, South Africa, GMN-BLN-ZMHU. O.d.: 7636. Ref.: 1802, 6681, 7716.]
- Xorides* (*Moerophora*) Förster, 1869. [Type-species: *Xylonomus rufipes* Gravenhorst, 1829. O.d.: 2318. Ref.: 6651, 149, 1802, 6682, 7990, 7713, 5701, 7706, 7682, 2795, 2780, 2475.]
- Macrosterotrichus* Kokujev, 1903. (*Xylonomus* subg.) [Type-species: *Xylonomus jakovlevi* Kokujev, 1903. O.d.: 4148. Ref.: 7990, 1394, 363, 1542.]
- austriacus* Clément, 1938 (*Macrosterotrichus*). Palearctic. [Type: ♀, GMN-BLN-ZMHU. O.d.: 1394. Ref.: 363, 5488, 257, 1542, 3946, 3544.]
- bischoffi* Clément, 1938 (*Macrosterotrichus*). Palearctic. [Type: ♀, GMN-BLN-ZMHU. O.d.: 1394. Ref.: 363, 5488, 257, 1542, 3946, 3544.]
- ephiatoides* Kriechbaumer, 1882 (*Xylonomus*). Palearctic. [Type: ♀, Germany, GMN-MCN-ZS. O.d.: 4324. Ref.: 1802, 6681, 6683, 4116, 4119, 5012, 6622, 1394, 192, 2864, 197, 363, 7706, 257, 6821, 1542, 3946, 4207, 6825, 4223.]
- hiatus* Wang & Gupta, 1995 (*Xorides*). Oriental. [Type: ♀, China, CNA-BJG-ASIZ. O.d.: 8232.]
- immaculatus* Cushman, 1933 (*Xorides*). Oriental. Palearctic. [Type: ♀, China, GMN-EBW-DEI. O.d.: 1744. Ref.: 7843, 7713, 7719, 2795, 1279, 1337, 2780, 8228, 4558, 3058.]
- indicus* Gupta & Chandra, 1972. Oriental. Preoccupied by *Xorides indicus* Szépligeti, 1914. [Type: ♀, India, USA-GVL-GUPTA. O.d.: 2795. Ref.: 2780.]
- investigator* Smith, 1874 (*Xylonomus*). Palearctic. [Type: ♀, Japan, UKD-LDN-NHM. O.d.: 7082. Ref.: 1802, 6681, 7775, 7784, 4925, 2147, 7818, 7843, 3667, 7713, 3671, 5155, 7706.]
- jakovlevi* Kokujev, 1903 (*Xylonomus* (*Macrosterotrichus*)). Palearctic. [Type: Russia, RSA-SPT-ZI. O.d.: 4148. Ref.: 5012, 1394, 7713, 7706, 257, 1542, 3946.]
- jezoensis* Matsumura, 1912 (*Xylonomus*). Palearctic. [Type: ♀, Japan, JPN-SPR-HU. O.d.: 4922. Ref.: 7775, 4925, 7713, 7706.]
- lineatus* Matsumura, 1912 (*Xylonomus*). [Type: ♀, Japan, JPN-SPR-HU. O.d.: 4922. Ref.: 7775, 7842, 7706.]
- lissopunctus* Gupta & Chandra, 1972. Oriental. [Type: ♀, India, USA-GVL-GUPTA. O.d.: 2795. Ref.: 2780.]
- nasensis* Uchida, 1956 (*Xorides*). Oriental. [Type: ♀, Japan, JPN-SPR-HU. O.d.: 7843. Ref.: 7719, 5134, 4495, 2780.]
- rufipes* Gravenhorst, 1829 (*Xylonomus*). Palearctic. [O.d.: 2636. Ref.: 1696, 4004, 892, 6651, 1802, 6680, 6681, 2476, 4116, 3251, 4117, 3146, 4119,

2922, 5012, 3098, 1394, 2849, 1223, 6804, 363, 7713, 219, 1501, 257, 1542, 4011, 3946, 2462, 3090, 4018, 6825, 4213, 4021, 4223, 4225.]

Xorides (*Xorides*) Latreille, 1809. [Type-species: *Ichneumon indicatorius* Latreille, 1806. O.d.: 4631. Ref.: 4632, 2630, 2636, 5356, 8328, 951, 6172, 3410, 3412, 7374, 7377, 2318, 7493, 1664, 6651, 149, 1802, 6681, 6682, 5179, 6683, 7990, 6271, 1208, 7775, 5012, 1394, 7648, 7710, 7655, 3441, 363, 7713, 382, 5488, 385, 7706, 7715, 242, 7682, 257, 7716, 2796, 5949, 1542, 1165, 2447, 2780, 2475.] Misidentified *Xorides* of authors = *Deuterororides* Viereck, 1914 or *Neoxorides* Clément, 1938.

Xylonomus Gravenhorst, 1829. (*Xorides* subg.) [Type-species: *Ichneumon irrigator* Fabricius, 1793. O.d.: 2636. Ref.: 5356, 8328, 6168, 951, 3410, 3412, 7374, 7377, 2318, 6042, 7493, 1664, 6651, 149, 1802, 6680, 6681, 6682, 5179, 5199, 7990, 1208, 7775, 5012, 1394, 363, 1542.]

Rhadina Förster, 1869. Preoccupied by *Rhadina* Billberg, 1828 (Aves). [Type-species: *Xylonomus ater* Gravenhorst, 1829. O.d.: 2318. Ref.: 6651, 149, 1802, 6682, 6771, 7990, 5701.]

Sichella Förster, 1869. [Type-species: *Xylonomus filiformis* Gravenhorst, 1829. O.d.: 2318. Ref.: 6651, 149, 1802, 6682, 7990, 1394, 363, 382, 5701, 385, 1542.]

Sterotrichus Förster, 1869. [Type-species: *Xylonomus pilicornis* Gravenhorst, 1829. O.d.: 2318. Ref.: 6651, 149, 1802, 6682, 7990, 7998, 5701.]

Perissocerus Smith, 1877. Preoccupied by *Perissocerus* Gerstaecker, 1868 (Diptera). [Type-species: *Perissocerus plumicornis* Smith, 1877. O.d.: 7085. Ref.: 1802, 6769, 7990.]

Pyramirhyssa Mocsáry, 1905. [Type-species: *Pyramirhyssa magnifica* Mocsáry, 1905. O.d.: 5081. Ref.: 6681, 7990, 7713, 7715, 7682, 2475.]

Periceros Schulz, 1906. [New name for *Perissocerus* Smith, 1877. O.d.: 6769. Ref.: 7713, 7715, 7682, 2475.]

Rhadinopimpla Schulz, 1911. [New name for *Rhadina* Förster, 1869. O.d.: 6771. Ref.: 1394, 363, 1542.]

Neoxylonomus Szépligeti, 1914. [Type-species: *Neoxylonomus antennalis* Szépligeti, 1914. O.d.: 7348. Ref.: 7682.]

Neoxylonomus Clément, 1938. Preoccupied by *Neoxylonomus* Szépligeti, 1914 (Ichneumonidae). [Type-species: *Neoxylonomus strandi* Clément, 1938. O.d.: 1394. Ref.: 363.]

Xylonominus Clément, 1938. [Type-species: *Xylonomus gracilicornis* Gravenhorst, 1829. O.d.: 1394. Ref.: 363, 1542.]

Amphixylonomus Benoît, 1952. (*Xylonomus* subg.) [Type-species: *Xylonomus kondiensis* Benoît, 1952. O.d.: 510.]

albopictus Cresson, 1870 (*Xylonomus*). Nearctic. [Type: ♀, U.S.A., USA-PLD-ANSP. O.d.: 1652. Ref.: 6040, 6064, 6068, 1664, 2991, 147, 1802, 6681, 1666, 7992, 6271, 1735, 5657, 7648, 2189, 7710, 7713, 1165.]

canadensis Provancher, 1875. [Type: ♀, Canada, CND-SFY-UL. O.d.: 6038. Ref.: 1664, 1802, 7648, 404.]

duplicatus Rohwer, 1913 (*Xylonomus* (*Moerophora*)). [Type: ♂, U.S.A., USA-WST-USNM. O.d.: 6265. Ref.: 6271, 7713.]

alpestris Habermehl, 1903 (*Xylonomus*). Palearctic. [Type: ♀, Germany. O.d.: 2869. Ref.: 2871, 6680, 6681, 5792, 7348, 5012, 1394, 3262, 3263, 3184, 3189, 3193, 6804, 363, 5488, 6814, 257, 1542, 3946, 6648, 3090, 6825, 4223.]

koromevi Kokujev, 1927 (*Sichella*). [Type: ♂, Russia, RSA-SPT-ZI. O.d.: 4169. Ref.: 5012, 1394, 7706, 257.]

annulatus Gravenhorst, 1829 (*Xylonomus*). Palearctic. [O.d.: 2636. Ref.: 1696, 1802, 6680, 6681, 5012, 1394, 363, 257, 1542, 3946, 4020.]

anthracinus Gupta & Chandra, 1974. Oriental. [Type: ♀, India, USA-GVL-GUPTA. O.d.: 2796. Ref.: 2780.]

armidae Gauld, 1997. Neotropical. [Type: ♀, Costa Rica, CTR-STD-INBC. O.d.: 2475.]

asperus Wang & Gupta, 1995. Oriental. [Type: ♀, China, CNA-BJG-ASIZ. O.d.: 8232.]

ater Gravenhorst, 1829 (*Xylonomus*). Palearctic. [Type: ♀, PLD-WCW-IZU. O.d.: 2636. Ref.: 1696, 6170, 3412, 7374, 3700, 8075, 8386, 8392, 6651, 4341, 7242, 1802, 6680, 6681, 2476, 7167, 7888, 3251, 5012, 5013, 440, 6622, 3098, 1394, 3263, 2841, 3184, 31, 6804, 363, 3817, 3822, 7706, 6815, 257, 1514, 1518, 1542, 4011, 3946, 6564, 3090, 4018, 6825, 7874.] Infrasubspecific names: *nigripes* (3251), *nigrofemorata* (3251).

var. *brunnescens* Clément, 1938 (*Rhadinopimpla ater* var.). [Type: ♀, GMN-MCN-ZS. O.d.: 1394. Ref.: 3544.]

var. *nigricoxis* Clément, 1938 (*Rhadinopimpla ater* var.). [Type: ♂, GMN-MCN-ZS. O.d.: 1394. Ref.: 3544.]

var. *rufosignatus* Clément, 1938 (*Rhadinopimpla ater* var. *rufosignatus*). [Type: ♀, GMN-MCN-ZS. O.d.: 1394. Ref.: 3544.]

atrox Townes, 1960. Nearctic. [Type: ♀, Canada, USA-SPL-UM. O.d.: 7713. Ref.: 1165.]

baueri Clément, 1938 (*Rhadinopimpla*). Palearctic. Preoccupied by *Xorides baueri* (Haupt, 1917). [Type: ♀, GMN-MCN-ZS. O.d.: 1394. Ref.: 6804, 206, 363, 242, 257, 1542, 3946, 3512, 6825, 3544.]

berlandi Clément, 1938. Palearctic. [Type: ♂, FCE-PRS-MNHN. O.d.: 1394. Ref.: 192, 221, 257, 1542, 3946, 4213, 3544, 4223.]

boharti Townes, 1960. Nearctic. [Type: ♂, U.S.A., USA-WST-USNM. O.d.: 7713. Ref.: 1165.]

brachylabis Kriechbaumer, 1889 (*Xylonomus*). Palearctic. [Type: ♀, Germany, GMN-MCN-ZS. O.d.: 4341. Ref.: 4612, 1802, 6680, 6681, 7887, 4127, 7888, 5792, 7127, 1174, 6873, 6877, 2652, 7027, 1347, 4121, 6622, 3098, 1394, 4130, 3271, 5472, 3184, 3115, 3189, 7196, 31, 6804, 363, 5101, 6814, 7706, 6815, 257, 1542, 3946, 2462, 6564, 3090, 6825, 7874, 4213, 7880, 4021, 4223.]

sachalinensis Uchida, 1928 (*Xylonomus*). [Type: ♀, Russia, JPN-SPR-HU. O.d.: 7775. Ref.: 7706.]

calidus Provancher, 1886 (*Xylonomus*). Nearctic. [Type: ♀, Canada, CND-SFY-UL. O.d.: 6072. Ref.: 1664, 2991, 1802, 6681, 2409, 6271, 1266, 5657, 7648, 2189, 7710, 7713, 1989, 404, 1165.]

pulcher Ashmead, 1890 (*Xylonomus*). [Type: ♀, U.S.A., USA-WST-USNM. O.d.: 126. Ref.: 2991,

- 1802, 6681.]
nasonii Davis, 1895 (*Aplomerus*). [Type: ♂, U.S.A., USA-PLD-ANSP. O.d.: 1833. Ref.: 1802, 5341, 6681, 1667.]
- californicus* Cresson, 1879 (*Xylonomus*). Nearctic. [Type: ♀, U.S.A., USA-PLD-ANSP. O.d.: 1661. Ref.: 1664, 2991, 1802, 6681, 1666, 6271, 7648, 7710, 7713, 1165.]
- catomus* Davis, 1897 (*Xylonomus*). [Type: ♀, U.S.A., USA-PLD-ANSP. O.d.: 1836. Ref.: 1802, 6681, 6271, 1667, 7648.]
- centromaculatus* Cushman, 1933. Oriental. [Type: ♀, China, GMN-EBW-DEI. O.d.: 1744. Ref.: 7719, 2796, 1279, 1337, 2780, 8232.]
- cerbonei* Porter, 1978 (*Xorides* (*Periceros*)). Neotropical. [Type: ♂, Mexico, MXC-MXC-MHN. O.d.: 5953.]
- cincticornis* Cresson, 1865 (*Xylonomus*). Nearctic. [Type: ♀, U.S.A., USA-PLD-ANSP. O.d.: 1645. Ref.: 1652, 1664, 2991, 1802, 6681, 1666, 6271, 7648, 7710, 7713, 1920, 1165.]
- modestus* Rohwer, 1913 (*Xylonomus*). [Type: ♂, U.S.A., USA-WST-USNM. O.d.: 6265. Ref.: 6271, 7713.]
- ssp. *rufus* Townes, 1960 (*Xorides cincticornis* ssp.). Nearctic. Preoccupied by *Xorides rufus* Habermehl, 1918. [Type: ♂, U.S.A., USA-WST-USNM. O.d.: 7713. Ref.: 1165.]
- crassitibialis* Uchida, 1932 (*Xylonomus*). Palearctic. [Type: ♀, Japan, JPN-SPR-HU. O.d.: 7790. Ref.: 3667, 3671, 7706.]
- csikii* Clément, 1938. Palearctic. [Type: ♂, Hungary, HGR-BDP-TMA. O.d.: 1394. Ref.: 363, 238, 257, 1542, 3946, 2462, 6648, 3333, 3544.]
- depressus* Holmgren, 1860 (*Xylonomus*). Palearctic. [Type: ♀, Sweden, SWD-SHM-NR. O.d.: 3412. Ref.: 8390, 892, 900, 1802, 6681, 6683, 6877, 5012, 1394, 3263, 192, 1226, 5564, 6804, 197, 363, 1238, 231, 6814, 7706, 242, 257, 1542, 3946, 6648, 3090, 6825.]
- linearis* Clément, 1938 (*Rhadinopimpla*). [Type: ♀, ATA-WEN-NM. O.d.: 1394. Ref.: 363, 1542, 3544.]
- eastoni* Rohwer, 1913 (*Xylonomus* (*Moerophora*)). Nearctic. [Type: ♀, U.S.A., USA-WST-USNM. O.d.: 6265. Ref.: 6271, 7648, 7710, 7713, 1165.]
- erigentis* Wang & Gupta, 1995. Palearctic. [Type: ♀, China, CNA-BJG-ASIZ. O.d.: 8232.]
- euthrix* Porter, 1975 (*Xorides* (*Periceros*)). Neotropical. [Type: ♀, Argentina, AGT-TCM-UNT. O.d.: 5949. Ref.: 5953, 2612.]
- filiformis* Gravenhorst, 1829 (*Xylonomus*). Palearctic. [O.d.: 2636. Ref.: 1696, 6168, 3999, 3412, 7374, 3700, 8078, 7493, 2187, 6651, 1990, 3698, 1802, 2871, 6680, 6681, 2476, 4127, 3023, 2180, 5838, 4117, 3147, 4119, 4120, 1629, 5012, 3098, 3258, 1394, 2841, 2845, 2849, 2850, 6627, 4642, 5472, 1194, 459, 6804, 209, 363, 1486, 6813, 6814, 6815, 257, 1542, 4011, 3946, 5886, 6564, 6648, 2266, 3090, 4018, 6825, 7876, 4213, 7880, 7882, 4021, 2547, 4223, 4225.]
- habermehli* Kiss, 1926 (*Xylonomus filiformis* var.). [Type: Hungary. O.d.: 4119. Ref.: 1394, 1194, 1542.]
- bicolor* Clément, 1938 (*Sichelia filiformis* var.). [Type: ♂, GMN-MCN-ZS. O.d.: 1394. Ref.: 1542, 3544.]
- var. *obscuripes* Kiss, 1929 (*Xylonomus filiformis* var.). Palearctic. [Type: ♂, Hungary. O.d.: 4120.]
- formosulus* Kokujev, 1912 (*Xylonomus*). Palearctic. [Type: ♀, Turkmenistan. O.d.: 4164. Ref.: 5012.]
- frigidus* Cresson, 1870 (*Xylonomus*). Nearctic. [Type: ♀, Canada, USA-PLD-ANSP. O.d.: 1652. Ref.: 6023, 6064, 6068, 1664, 2991, 1802, 154, 6681, 1666, 6271, 1735, 7648, 7710, 7713, 1165.]
- plesius* Rohwer, 1913 (*Xylonomus*). [Type: ♀, USA-WST-USNM. O.d.: 6265. Ref.: 6271, 7713.]
- yukonensis* Rohwer, 1913 (*Xylonomus* (*Moerophora*)). [Type: ♀, U.S.A., USA-WST-USNM. O.d.: 6265. Ref.: 6271, 7713.]
- fuligator* Thunberg, 1822 (*Ichneumon*). Palearctic. [O.d.: 7537. Ref.: 7538, 1802, 6310, 3251, 5012, 5791, 3098, 1394, 3263, 4640, 4130, 4642, 1194, 6804, 3133, 363, 221, 219, 4062, 5863, 257, 258, 1542, 5874, 3946, 2462, 6564, 6648, 4186, 3090, 6825, 7874, 4213, 4020, 4223, 4225.]
- sordator* Thunberg, 1822 (*Ichneumon*). [Type: ♂, Sweden, SWD-USL-UU. O.d.: 7537. Ref.: 7538, 1802, 6310.]
- pilicornis* Gravenhorst, 1829 (*Xylonomus*). [O.d.: 2636. Ref.: 1689, 1696, 1913, 3412, 7374, 7377, 4861, 8078, 7493, 892, 6651, 1990, 4612, 7242, 1802, 7282, 2871, 699, 6681, 2476, 5179, 6683, 7887, 5431, 6437, 7127, 1174, 2967, 4119, 7628, 1441, 2943, 1629, 2849, 31.]
- gracilicornis* Gravenhorst, 1829 (*Xylonomus*). Palearctic. [O.d.: 2636. Ref.: 1696, 3412, 7374, 3700, 6651, 1802, 6680, 6681, 4614, 7348, 4116, 2887, 687, 4987, 1208, 4119, 7775, 1629, 5012, 3098, 3258, 1394, 5471, 5472, 1223, 192, 6804, 363, 7706, 242, 1504, 251, 257, 1542, 3946, 5886, 6648, 4186, 6825, 7877, 4213, 2547, 4223, 4225.]
- baueri* Haupt, 1917 (*Xylonomus*). [Type: ♀, Germany, GMN-MCN-ZS. O.d.: 3023. Ref.: 3512.]
- var. *atratus* Obertl, 1946 (*Xylonomus gracilicornis* var.). Palearctic. [Type: Czechoslovakia. O.d.: 5471. Ref.: 363.]
- var. *nigripes* Kiss, 1929 (*Xylonomus gracilicornis* var.). Palearctic. [Type: ♀, Hungary, HGR-BDP-TMA. O.d.: 4120. Ref.: 3544.]
- nigripes* Clément, 1938 (*Xylonomus gracilicornis* var.). Preoccupied by *Xorides nigripes* (Kiss, 1929). [Type: ♀, Hungary, HGR-BDP-TMA. O.d.: 1394. Ref.: 6681, 363, 3544.]
- gravenhorstii* Curtis, 1831 (*Xylonomus*). Palearctic. [Type: ♀, United Kingdom, ATL-MBE-NMV. O.d.: 1689. Ref.: 1696, 1913, 4861, 1802, 6681, 2258, 2462, 6825, 2547, 4223, 4225.]
- securicornis* Holmgren, 1860 (*Xylonomus*). [Type: ♀, Sweden. O.d.: 3412. Ref.: 8078, 1802, 6681, 5179, 6683, 6314, 6552, 5012, 3098, 1394, 4130, 4642, 5472, 3184, 3189, 459, 6804, 363, 231, 238, 242, 1501, 6815, 1505, 5862, 257, 1542, 7285, 3946, 6564, 6648, 320, 4186, 3090, 4208, 4213, 4020, 4021.]
- glyptus* Thomson, 1877 (*Xylonomus*). [Type: SWD-LND-ZI. O.d.: 7493. Ref.: 1802, 6681, 2263.]
- clavicornis* Kriechbaumer, 1879 (*Xylonomus*). [O.d.: 4302. Ref.: 1802, 91, 6681, 6683, 5012, 1223.]
- distinguendus* Magretti, 1884 (*Xylonomus*). [O.d.: 4796. Ref.: 1802, 6681.]
- hungaricus* Szépligeti, 1899 (*Sichelia hungarica*).

- [Type: ♀, Hungary. O.d.: 7334. Ref.: 1802, 6681, 7348, 4119.]
- seticornis* Strobl, 1902 (*Xylonomus*). [Type: ♀, Yugoslavia. O.d.: 7282. Ref.: 6680, 6681, 6683, 3023, 4117, 2652, 4121, 5012, 6627, 7285.]
- rufoscutellatus* Habermehl, 1918 (*Xylonomus seticornis* f. *rufoscutellatus*). [Type: ♀, Algeria. O.d.: 2887. Ref.: 1394, 3514.]
- kokujevi* Meyer, 1922 (*Xylonomus*). [Type: Russia. O.d.: 4986. Lapse: kokujevi. Ref.: 5012, 1394.]
- rufus* Kiss, 1924 (*Xylonomus pilicornis* var.). [Type: ♂, Hungary. O.d.: 4117.]
- caucasicus* Shestakov, 1925 (*Xylonomus*). [O.d.: 6957. Ref.: 1394.]
- romani* Clément, 1938. [Type: ♀, SWD-SHM-NR. O.d.: 1394. Ref.: 1542, 3544.]
- harringtoni* Rohwer, 1920. Nearctic. [New name for *Xylonomus canadensis* Harrington, 1891. O.d.: 6271. Ref.: 7710, 7713, 1165.]
- canadensis* Harrington, 1891 (*Xylonomus*). Invalidated by pre-1961 replacement. Preoccupied by *Xorides canadensis* Provancher, 1875. [Type: USA-WST-USNM. O.d.: 2991. Ref.: 1802, 6681, 6271.]
- hedwigi* Clément, 1938. Palearctic. [Type: ♂, GMN-BLN-ZMHU. O.d.: 1394. Ref.: 3098, 7196, 6804, 363, 257, 1542, 3946, 6825, 4020, 3544, 4021.]
- hingganensis* Wang & Gupta, 1995. Palearctic. [Type: ♀, China, CNA-BJG-ASIZ. O.d.: 8232.]
- humos* Gauld, 1997. Neotropical. [Type: ♀, Costa Rica, CTR-STD-INBC. O.d.: 2475.]
- idunae* Gauld, 1997. Neotropical. [Type: ♀, Costa Rica, CTR-STD-INBC. O.d.: 2475.]
- indicatorius* Latreille, 1806 (*Ichneumon*). Palearctic. [O.d.: 4630. Ref.: 4631, 1802, 6681, 3098, 1394, 6804, 197, 363, 5862, 257, 1542, 4011, 3946, 3090, 6825, 7880, 4020, 4223.]
- ferrugatus* Gravenhorst, 1829 (*Xylonomus*). [O.d.: 2636. Ref.: 1696, 3412, 7493, 1802, 6680, 6681, 2476, 4986, 3146, 1440, 7628, 2927, 5012, 2849, 1224, 1227.]
- quadrimaculatus* Clément, 1938 (*indicatorius* var.). [Type: ♀, Germany, GMN-MCN-ZS. O.d.: 1394. Ref.: 363, 1542, 3512, 3544.]
- insularis* Cresson, 1879 (*Poemenia*). Nearctic. [Type: ♀, Canada, USA-PLD-ANSP. O.d.: 1661. Ref.: 1664, 2991, 1802, 6681, 1666, 6271, 7648, 7277, 2732, 7710, 7713, 6984, 1165.]
- irrigator* Fabricius, 1793 (*Ichneumon*). Palearctic. [Type: ♀, Germany, DMK-CPH-UZM. O.d.: 2169. Ref.: 5599, 2174, 5607, 2628, 2632, 7537, 2636, 1696, 8481, 6168, 3412, 7374, 7377, 3700, 4861, 8386, 7493, 6651, 3698, 1802, 7282, 2871, 699, 6681, 2476, 5179, 6683, 4127, 5792, 7348, 3251, 6437, 7127, 1174, 1208, 4119, 1347, 1629, 5012, 5013, 5791, 440, 6622, 3098, 1394, 3262, 3263, 2849, 4130, 4642, 1194, 2586, 1223, 7196, 31, 2864, 6804, 363, 210, 5101, 8498, 1502, 7706, 242, 1501, 4062, 7682, 257, 1542, 8468, 4011, 3946, 2462, 6564, 6568, 3090, 4018, 6825, 7365, 7874, 4213, 7880, 7882, 4020, 4021, 2547, 4223.]
- Infrasubspecific name: *nigriventris* (3262).
- planus* Sedivy, 1958. [Type: ♀, Czechoslovakia, CCA-PRH-NM. O.d.: 6804.]
- iwatensis* Uchida, 1928 (*Xylonomus*). Palearctic. [Type: ♀, Japan, JPN-SPR-HU. O.d.: 7775. Ref.: 7706.]
- kondensis* Benoit, 1952 (*Xylonomus (Amphixylonomus)*). Ethiopian. [Type: ♀, Zaire, BGM-TVIR-MRAC. O.d.: 510. Ref.: 7716.]
- konumensis* Uchida, 1928 (*Xylonomus*). Palearctic. [Type: ♀, Russia, JPN-SPR-HU. O.d.: 7775. Ref.: 7813, 5101, 7706.]
- magnificus* Mocsáry, 1905 (*Pyramirhyssa magnifica*). Neotropical. [Type: ♀, Brazil, HGR-BDP-TMA. O.d.: 5081. Ref.: 6681, 7715, 5949, 5953, 2612.]
- maudae* Davis, 1895 (*Xylonomus*). Nearctic. [Type: ♀, U.S.A., USA-PLD-ANSP. O.d.: 1833. Ref.: 1802, 6681, 6271, 1667, 7648, 7710, 7713, 1165.]
- medius* Townes, 1960. Nearctic. [Type: ♀, U.S.A., USA-WST-USNM. O.d.: 7713. Ref.: 1165.]
- minimus* Gupta & Chandra, 1974. Oriental. [Type: ♀, India, USA-GVL-GUPTA. O.d.: 2796. Ref.: 2780.]
- minutus* Clément, 1938. Palearctic. [Type: ♀, FCE-PRS-MNH. O.d.: 1394. Ref.: 3110, 192, 257, 1542, 3946, 5886, 6825, 4213, 3544, 4223.]
- neoclyti* Rohwer, 1915 (*Moerophora*). Nearctic. [Type: ♂, U.S.A., USA-WST-USNM. O.d.: 6268. Ref.: 6271, 7648, 7710, 7713, 1165.]
- niger* Pfeffer, 1913 (*Xylonomus*). Palearctic. [Type: ♀, Germany. O.d.: 5792. Ref.: 1394, 363, 6813, 6815, 257, 1542, 3946, 2462, 5886, 6825, 7874, 7877, 4021.]
- bicolor* Clément, 1938 (*niger* var.). [Type: ♀, GMN-MCN-ZS. O.d.: 1394. Ref.: 5472, 3184, 363, 5862, 1542, 5886, 7877, 3544.]
- var. *rufus* Sedivy, 1958 (*Xorides niger* var. *rufa*). Palearctic. Preoccupied by *Xorides rufus* Habermehl, 1918. [O.d.: 6804. Ref.: 363.]
- nigristomus* Gupta & Chandra, 1974. Oriental. [Type: ♀, India, USA-GVL-GUPTA. O.d.: 2796. Ref.: 2780.]
- ornatus* Tosquinet, 1903 (*Moansa*). Oriental. [Type: ♀, Indonesia, IITL-GNA-MCSN. O.d.: 7639. Ref.: 6681, 7713, 7719, 2796, 2780.]
- peniculus* Townes, 1960. Nearctic. [Type: ♀, U.S.A., USA-WST-USNM. O.d.: 7713. Ref.: 1165.]
- pictus* Townes, 1960. Nearctic. [Type: ♀, U.S.A., USA-GVL-AEI. O.d.: 7713. Ref.: 1165.]
- planus* Townes, 1960. Nearctic. Preoccupied by *Xorides planus* Sedivy, 1958. [Type: ♀, U.S.A., USA-ITC-CU. O.d.: 7713. Ref.: 1165.]
- plumicornis* Smith, 1877 (*Perissocerus*). Neotropical. [Type: ♀, Brazil, UKD-LDN-NHM. O.d.: 7085. Ref.: 1802, 6769, 7715, 5953, 2612.]
- antennalis* Szépligeti, 1914 (*Neoxylonomus*). [Type: ♀, Brazil, HGR-BDP-TMA. O.d.: 7348. Ref.: 7715.]
- praecatorius* Fabricius, 1793 (*Ichneumon*). Palearctic. [O.d.: 2169. Ref.: 2174, 4629, 2632, 7537, 2636, 7726, 1696, 6168, 3412, 7374, 7377, 4861, 7493, 892, 6651, 908, 1990, 1802, 7282, 2871, 699, 6680, 6681, 2476, 5179, 2047, 4127, 5792, 7123, 7127, 4986, 3146, 4119, 2652, 1441, 4120, 2943, 1629, 5012, 5013, 3098, 1394, 5083, 2001, 2841, 490, 2849, 4130, 5472, 1194, 7196, 459, 2864, 6804, 363, 1478, 8498, 6814, 242, 6815, 257, 258, 1542, 6984, 3946, 2462, 6563, 5886, 6564, 6648, 6568, 3090, 184, 3333, 6825, 7365, 7874, 7876, 7878, 4213, 7880, 7881, 4020, 4021, 2547, 4223, 4225.]
- ? *articulatus* Geoffroy, 1785 (*Ichneumon*). [O.d.: 2325. Ref.: 1802, 257.]

- ? *falsatorius* Olivier, 1792 (*Ichneumon*). [Emendation for *Ichneumon articulatus* Geoffroy, 1785. O.d.: 5512. Ref.: 1802, 257.]
- parvulus* Gravenhorst, 1829 (*Xylonomus*). [O.d.: 2636. Ref.: 1696, 1913, 1802.]
- praecatorius* Marshall, 1872 (*Xylonomus*). Preoccupied by *Xorides praecatorius* (Fabricius, 1793). [Emendation for *Ichneumon praecatorius* Fabricius, 1793. O.d.: 4861.]
- rufopictus* Kiss, 1926 (*Xylonomus praecatorius* var. *rufopicta*). [Type: ♂, Hungary. O.d.: 4119.]
- ruficoxis* Kiss, 1929 (*Xylonomus praecatorius* var.). Preoccupied by *Xylonomus ruficoxis* Rohwer, 1913. [Type: ♂, Hungary. O.d.: 4120.]
- temporalis* Kiss, 1929 (*Xylonomus praecatorius* var.). [Type: ♂, Hungary. O.d.: 4120. Ref.: 1501.]
- holsaticus* Heinrich, 1951. [Type: Germany. O.d.: 3191.]
- var. *funebis* Kiss, 1929 (*Xylonomus praecatorius* var.). Palearctic. [Type: ♀, Hungary. O.d.: 4120.]
- ssp. *rufigaster* Constantineanu, Ciochia, Constantineanu & Mustata, 1967 (*Xorides praecatorius* ssp.). Palearctic. [Type: ♀, Romania. O.d.: 1501.]
- rileyi* Ashmead, 1890 (*Xylonomus*). Nearctic. [Type: ♀, U.S.A., USA-WST-USNM. O.d.: 126. Ref.: 2991, 1802, 6681, 6271, 7648, 2189, 7710, 6978, 7713, 6984, 1165.]
- lepturae* Rohwer, 1913 (*Xylonomus*). [Type: ♀, U.S.A., USA-WST-USNM. O.d.: 6266. Ref.: 6271, 7713.]
- ruficoxis* Rohwer, 1913 (*Xylonomus*). [Type: ♀, U.S.A., USA-WST-USNM. O.d.: 6265. Ref.: 6271, 7648, 2189, 7710, 7713.]
- rudis* Townes, 1960. Nearctic. [Type: ♀, U.S.A., USA-LRC-UK. O.d.: 7713. Ref.: 1165.]
- rusticus* Desvignes, 1856 (*Xylonomus*). Palearctic. [Type: ♀, United Kingdom, UKD-LDN-NHM. O.d.: 1913. Ref.: 4861, 1802, 6681, 5179, 4130, 2258, 2462.]
- sapporensis* Uchida, 1928 (*Xylonomus*). Palearctic. [Type: ♀, Japan, JPN-SPR-HU. O.d.: 7775. Ref.: 4925, 2147, 7706.]
- nigritibiae* Uchida, 1952 (*Xylonomus sapporensis* var.). [Type: ♀, Japan, JPN-SPR-HU. O.d.: 7835. Ref.: 3667, 3671, 7706.]
- secos* Gauld, 1997. Neotropical. [Type: ♀, Costa Rica, CTR-STD-INBC. O.d.: 2475.]
- sejugatus* Brues, 1910 (*Xylonomus*). Fossil. [Type: ♀, U.S.A., USA-CBG-MCZ. O.d.: 935.]
- semirufus* Townes, 1960. Nearctic. [Type: ♀, U.S.A., USA-GVL-AEI. O.d.: 7713. Ref.: 1165.]
- sepulchralis* Holmgren, 1860 (*Xylonomus*). Palearctic. [Type: ♀, Sweden, SWD-SHM-NR. O.d.: 3412. Ref.: 8392, 1802, 6680, 6681, 5792, 6328, 4119, 7775, 5001, 4925, 5012, 3258, 1394, 3262, 363, 5101, 7581, 7706, 238, 257, 1514, 1542, 3946, 3090, 6825, 4213, 2547, 4223.]
- moiwasanus* Matsumura, 1912 (*Xylonomus*). [Type: ♀, Japan, JPN-SPR-HU. O.d.: 4922. Ref.: 7706.]
- schmiedeknechti* Clément, 1938 (*Sichelia sepulchralis* var.). [Type: ♀, GMN-BLN-ZMHU. O.d.: 1394. Ref.: 3262, 3263, 3110, 3114, 363, 1542, 6648, 3544.]
- shevyrevi* Meyer, 1926 (*Xylonomus*). Palearctic. [Type: ♀, Azerbaijan. O.d.: 4993. Ref.: 5012.]
- stepposus* Kasparyan, 1981. Palearctic. [Type: ♀, Russia, RSA-SPT-ZI. O.d.: 3946.]
- stigmapterus* Say, 1824 (*Acaenitus*). Nearctic. [Type: ♀, Canada, USA-PLD-ANSP. O.d.: 6582. Ref.: 1652, 6023, 6064, 6068, 2991, 7042, 147, 1802, 5341, 6681, 2393, 7992, 6271, 1778, 1735, 5994, 7648, 7277, 2189, 7710, 7713, 7664, 1165.]
- tartarus* Morley, 1913 (*Xylonomus*). [Type: ♀, U.S.A., UKD-OFD-HOPE. O.d.: 5199. Ref.: 5318, 7663, 7664.]
- ssp. *floridanus* Ashmead, 1890 (*Xylonomus*). Nearctic. [Type: ♀, U.S.A., USA-WST-USNM. O.d.: 126. Ref.: 2991, 1802, 6681, 6271, 7648, 2189, 7710, 7713, 1165.]
- strandii* Clément, 1938 (*Neoxylonomus*). Palearctic. [Type: ♀, Germany, GMN-BLN-ZMHU. O.d.: 1394. Ref.: 363, 7719, 5488, 257, 3946, 3544.]
- syrinx* Gauld, 1997. Neotropical. [Type: ♀, Costa Rica, CTR-STD-INBC. O.d.: 2475.]
- tamora* Gauld, 1997. Neotropical. [Type: ♀, Costa Rica, UKD-LDN-NHM. O.d.: 2475.]
- thanatos* Gauld, 1997. Neotropical. [Type: ♀, Costa Rica, UKD-LDN-NHM. O.d.: 2475.]
- ? *tornatus* Schiødtte, 1839 (*Xylonomus*). Palearctic. [Type: ♀, Denmark. O.d.: 6633.]
- townesi* Baltazar, 1961 (*Xorides (Sichelia)*). Oriental. [Type: ♀, Philippines, USA-GVL-AEI. O.d.: 382. Ref.: 388, 2780.]
- vitiosus* Turner, 1919 (*Xylonomus*). Oriental. [Type: ♀, Laos, UKD-LDN-NHM. O.d.: 7748. Ref.: 7713, 7719, 2796, 2780.]
- wenzeli* Gauld, 1997. Neotropical. [Type: ♀, Costa Rica, USA-GVL-AEI. O.d.: 2475.]
- xanthisma* Porter, 1975. Neotropical. [Type: ♀, Argentina, AGT-TCM-UNT. O.d.: 5949. Ref.: 5953, 2612.]
- yamai* Gauld, 1997. Neotropical. [Type: ♀, Costa Rica, CTR-STD-INBC. O.d.: 2475.]